

North Carolina State Health Director's Standing Order for Nicotine Replacement Therapy March 24, 2022

Pursuant to S.L. 2021-110, this standing order signed by the North Carolina State Health Director authorizes immunizing pharmacists practicing in the state of North Carolina and licensed by the North Carolina Board of Pharmacy to dispense, deliver, or administer the following nicotine replacement therapy products as directed below. https://www.ahrq.gov/prevention/guidelines/tobacco/clinicians/index.html

Nicotine Replacement Therapy (NRT) Dispensing Protocol

Nicotine Replacement Therapy (NRT) Dispensing Protocol				
Eligible Candidates				
	This standing order may be used for persons < 18 years of age with a parent or legal guardian consent.			
Initiate therapy	High Nicotine Use*^	Medium Nicotine Use*^	Low Nicotine Use*^	
based on maximum	11+ cigarettes per day OR ≥2 cans or	5-10 cigarettes per day OR	0-4 cigarettes per day OR	
use of nicotine/ day	pouches per week of snuff or chew OR 6-	1 to 2 cans or pouches per week of	less than 1 can or pouch	
	12+mg/mL e-liquid	snuff or chew OR 3-6mg/mL e-liquid	per week of snuff or chew	
at therapy initiation				
*Combination	Per Product Label:	Per Product Label:	Per Product Label:	
Nicotine	•Nicotine Patch 21mg/24hrs for 8 weeks.	•Nicotine Patch 14mg/24hrs for 8	Nicotine Gum 2mg every	
Replacement therapy	Then,	weeks. Then,	hour as needed for	
is strongly	•Nicotine Patch 14mg/24hrs for 2 weeks.	Nicotine Patch 7mg/24hrs for 4	cravings. (Max	
recommended.	Then,	weeks.	20pieces/day) x 12 weeks.	
Monotherapy may	Nicotine Patch 7mg/24hrs for 2			
also be appropriate.	weeks.			
*Therapy choice				
should be based on	AND / OR any of the following as needed	AND / OR any of the following as	<u>OR</u>	
time to first use,	NRT products	needed NRT products	Nicotine lozenge 2mg	
quantity, patient	 Nicotine Gum 4mg every hour as needed 	Nicotine Gum 2mg every hour as	every hour as needed for	
preference and	for cravings. (Max 20pieces/day) x 12	needed for cravings. (Max 20	cravings. (Max 15/day) x	
comorbidities, data	weeks.	pieces/day) x 12 weeks.	12 weeks.	
from past attempts,				
and desired quit date.	<u>OR</u>	<u>OR</u>	<u>OR</u>	
	 Nicotine lozenge 4mg every hour as 	 Nicotine lozenge 2mg every hour as 	Nicotine Oral Inhaler	
NRT use in women	needed for cravings. (Max 15/day) x 12	needed for cravings. (Max 15/day) x	Puff 6-8 cartridges per day	
who are pregnant or	weeks.	12 weeks.	as needed for cravings x	
breastfeeding: the			12 weeks.	
patient should be	<u>OR</u>	<u>OR</u>		
educated on the risks	Nicotine Oral Inhaler	Nicotine Oral Inhaler	<u>OR</u>	
of smoking or vaping versus the unknown	Puff 6-16 cartridges per day as needed for	Puff 6-16 cartridges per day as needed	Nicotine Nasal Inhaler	
risks of NRT. If the	cravings x12 weeks.	for cravings x12 weeks.	1-2 doses/hour; 8-20	
			doses per day as needed	
patient consents to NRT, then	<u>OR</u>	<u>OR</u>	for cravings x 12 weeks.	
*	Nicotine Nasal Inhaler	Nicotine Nasal Inhaler		
intermittent delivery formulations (gum,	1-2 doses/hour; 8-40 doses per day as	1-2 doses/hour; 8-40 doses per day as		
lozenge or inhaler)	needed for cravings x 12 weeks.	needed for cravings x 12 weeks.		
are believed to be				
safer than continuous				
delivery (avoid use of				
Transdermal Dermal				
patch). If the				
patient is pregnant,				
educate on				
importance of				
importance or				



PCP/ObGyn for				
further prenatal care.				
Patient Education &	Every person dispensed NRT pursuant to this standing order shall receive education regarding appropriate use and			
Behavioral Support	potential adverse effects for the provided NRT product(s). Patients shall also be provided with behavioral support			
	education or provided with referral information for support services. All North Carolinians can receive support for			
	quitting through QuitlineNC by calling 1-800-QUITNOW (1-800-784-8669), texting READY at 200-400 or through the			
	website www.QuitlineNC.com.			
Refills	PRN			
Contraindications	Per product labeling:			
*If patient has any of the	Myocardial infarction or Stroke/TIA within the last 2 weeks.			
following, refer to medical	Diagnosed with worsening or serious angina within the last 6 months			
provider for further care.	Diagnosed within last 6 months with very rapid or irregular heartbeat that required a change in activities or			
	addition of medication.			
	A history of known hypersensitivity or serious adverse reaction to NRT or any of its components.			
A	Contraindications for Nicotine Patch – Severe eczema or psoriasis Pharmacists choosing to dispense NRT under the authority of this standing order shall notify the patient's primary			
Notification of	care provider within 72 hours after administration. Notification should include the pharmacist's name and NPI #,			
primary care	and the pharmacy/practice name and phone number. If the patient does not have a primary care provider, the			
provider:	pharmacist shall counsel the patient regarding the benefits of establishing a relationship with a primary care			
	provider, and provide information regarding primary care providers, including private practices, federally qualified			
	health centers, free clinics, or local health departments serving the area in which the patient is located.			
	meanificenters, free clinics, or local health de	partificates serving the area in which the pa	atient is located.	
h . I				
(Tilson				
Approved by: Date signed: 3-24-2022				
Elizabeth Cuervo Tilson, MD, MPH				
Linzabeth Caci vo Tilboll, IVID	7) 1911 11			

NPI: 1760540421

(<u>Legal Authority Session Law 2021-110 HouseBill 96</u> This order is effective immediately upon signing and may be revised or revoided by the State Health Director according to his/her discretion. This order shall remain in effect until the later of the development of the protocols described in Section 4(a) Session Law 2021-110 HouseBill 96 or January 1, 2023.